

Analysis of Activation - Induced and Post - Activation Undershoot $\Delta R2^*$ and $\Delta R2$ Magnitudes and Ratios at 1.5 Tesla using Synchronous Gradient-Echo and Spin-Echo (SGS) - EPI

Peter A. Bandettini and Kathleen M. Donahue

Biophysics Research Institute, Medical College of Wisconsin, Milwaukee, WI, USA

Introduction:

This study involves the use of a synchronous gradient - echo and spin-echo (SGS) - echo planar imaging (EPI) sequence (1) to analyze, on a voxel - wise basis, relative gradient-echo and spin-echo changes during activation and during the post - activation undershoot. Simultaneous collection of gradient-echo and spin-echo time series provides precise spatial and temporal registration not achievable using separate time series.

The questions addressed on a voxel-wise basis in this study include: What is the correlation between the signal change magnitude and post activation undershoot magnitude? What is the correlation between $\Delta R2^* / \Delta R2$ ratios during activation and during the post - activation undershoot? What is the correlation between $\Delta R2^*$ and $\Delta R2$ ratios and magnitudes of each?

Methods:

SGS-EPI was performed using a local three axis gradient coil at 1.5T (GE Signa). Gradient-echo TE = 30 ms, and Spin-echo TE = 110 ms. TR = 1 sec. Two axial planes (voxel volume = $3.75 \times 3.75 \times 10 \text{ mm}^3$) were obtained: one containing visual cortex and one containing motor cortex. Subjects viewed a red 8 Hz flashing LED display (GRASSTTM goggles), and performed bilateral finger tapping when the visual stimulus was on. Each time series was 64 sec. in duration. Stimulus timing was 20 sec. off, 20 sec. on, and 24 sec. off. The time between successive series was at least 2 minutes. Thirty time series were collected, spatially registered, and averaged. Correlation analysis was performed to determine regions of motor and visual activation. From these regions, voxel-wise analysis was carried out.

Results and Conclusions:

Figure 1 shows simultaneously-acquired $\Delta R2^*$ and $\Delta R2$ time series from the motor and visual cortex of one subject. $\Delta R2^*(*) = -\ln(S_{\text{act}} / S_{\text{rest}}) / \text{TE}$. Visual cortex shows a larger undershoot than motor cortex. While this is typical, undershoots are still observable in motor cortex.

Figure 2 is a set of scatter plots from voxels in visual cortex of both subjects, showing $\Delta R2$ and $\Delta R2^*$ during activation (x axis) and during the post-activation undershoot (y-axis). $\Delta R2^*(*)$ for both the increase and undershoot appear correlated.

Figure 3 is a set of scatter plots showing a comparison of activation - induced $\Delta R2^*$ and $\Delta R2$ with $\Delta R2^* / \Delta R2$. The two plots show significantly different shapes. $\Delta R2^*$ shows a monotonic correlation with $\Delta R2^* / \Delta R2$ - suggesting that as $\Delta R2^*$ increases (likely with higher blood volume per voxel), the average vessel radius increases. The cluster of voxels showing the highest $\Delta R2^*$ shows a $\Delta R2^* / \Delta R2$ ratio of only 3 to 5. Based on susceptibility contrast models (2-4), this ratio corresponds to a radius of about 100 μm - neglecting intravascular perturbation effects. The largest fMRI signal changes are thought to arise from vessels larger than 100 μm - which should correspond to a considerably higher $\Delta R2^* / \Delta R2$ ratio. These results appear to indicate a significant contribution of red blood cell susceptibility perturbations - having a $\Delta R2^* / \Delta R2$ ratio of about 1.5 (5) - which factor in to reduce $\Delta R2^* / \Delta R2$ to the value measured.

The relationship between $\Delta R2^* / \Delta R2$ and $\Delta R2$ is not monotonic. The voxels showing the highest $\Delta R2$ also show a $\Delta R2^* / \Delta R2$ ratio of 3 to 5, which is a larger ratio than would correspond to capillary effects. This also implies an intravascular (red blood cell) contribution to $\Delta R2$.

Table 1 summarizes the results from visual and motor cortex voxels of each subject. The increase and undershoot $\Delta R2^* / \Delta R2$ ratios are not significantly different.




Figure 1: Simultaneously-acquired $\Delta R2^*$ and $\Delta R2$ time series from visual and motor cortex.




Figure 2: Voxel - wise scatter plots of the relation between the activation - induced signal change and the post - activation undershoot a. $\Delta R2^*$ and b. $\Delta R2$.




Figure 3: Voxel-wise scatter plots of the relation between $\Delta R2^* / \Delta R2$ ratio and a. $\Delta R2^*$ and b. $\Delta R2$ magnitudes.

	$\Delta R2^*$	$\Delta R2$	$\Delta R2^* / \Delta R2$	$\Delta R2^* / \Delta R2$	$\Delta R2^* / \Delta R2$
1 i V	$-.58 \pm .04$	$-.18 \pm .01$	$3.32 \pm .13$	$-3.6 \pm .9$	$-3.0 \pm .7$
2 i V	$-.45 \pm .03$	$-.12 \pm .01$	$4.13 \pm .35$	$-3.1 \pm .6$	$-2.4 \pm .3$
1 u V	$.13 \pm .01$	$.06 \pm .01$	$2.31 \pm .48$		
2 u V	$.15 \pm .01$	$.05 \pm .04$	$3.04 \pm .46$		
1 i M	$-.55 \pm .09$	$-.18 \pm .02$	$3.39 \pm .32$	-5.8 ± 4.9	$-4.1 \pm .3$
2 i M	$-.51 \pm .05$	$-.16 \pm .01$	$3.15 \pm .20$	-2.4 ± 2.5	-2.6 ± 1.1
1 u M	$.01 \pm .02$	$.02 \pm .01$	$1.48 \pm .80$		
2 u M	$.01 \pm .01$	$.02 \pm .01$	$.83 \pm .66$		

Table 1: Left three columns: Relaxation rates and ratios from two subjects (1 and 2) and corresponding to the signal increase (i) and undershoot (u) in the visual (V) and motor (M) cortex. Right two columns: $\Delta R2^*$ and $\Delta R2$ increase to undershoot ratios.

References:

1. P. A. Bandettini, et al. in "Proc., SMRM, 12th Annual Meeting, New York, 1993," p. 169.
2. S. Ogawa et al. *Biophysical J* 64, 803-812 (1993).
3. P. A. Bandettini, E. C. Wong, *Int. J. of Imag. Syst. and Tech.* 6, 134-152 (1995).
4. J. L. Boxerman, et al. *Magn. Reson. Med.* 34, 555-566 (1995).
5. B. E. Hoppel, et al. *Magn. Reson. Med.* 30, 715-723 (1993).

1998 Abstract Form for Scientific Presentations
INTERNATIONAL SOCIETY FOR
MAGNETIC RESONANCE IN MEDICINE
SIXTH SCIENTIFIC MEETING
SYDNEY, AUSTRALIA
APRIL 18-24, 1998

ABSTRACT DEADLINE:

Received no later than NOVEMBER 18, 1997.
All copyrights to accepted abstracts become the
property of the ISMRM. No proprietary information
may be withheld by authors.

The authors have submitted this abstract to ISMRM
in addition to their journal. I consent.

First name

Last name

☐ I prefer Oral Presentation but willing to present as a poster

☐ Video required (available only for Oral Presentations)

☐ I prefer Poster but willing to make Oral Presentation

☐ Poster only

☐ The authors certify that any work with human or animal subjects complies
with the guiding policies of their national or institutional regulatory bodies

Topic categories (please fill in both):

Enter "old" category number 0206

Enter new category number 2.2.2.1.12.16

Keywords fMRI RCU undershoot

SEE BACK OF FORM FOR ALL INSTRUCTIONS

Please type the name and complete mailing address of the first author

Name Peter A. Bandettini

Address Biophysics Res. Inst., Medical
College of WI, 8701 Watertown Plank
Rd., Milwaukee, WI 53226

Country U.S.A.

Telephone (414) 456-6036

FAX (414) 456-6512

e-mail pab@mcw.edu